arXiv:cond-mat/0207676v2 [cond-mat.stat-mech] 12 Aug 2002

Number-conserving rate equation for sympathetic cooling of a boson gas

A. N. Salgueiro[1]*
[1] *Max-Planck Institut für Physik komplexer Systeme, Nöthnitzer Strasse 38, D-01187, Dresden, Germany*

(February 1, 2008)

We derive a particle number-conserving rate equation for the ground state and for the elementary excitations of a bosonic system which is in contact with a gas of a different species (sympathetic cooling). We use the Giradeau-Arnowitt method and the model derived by Lewenstein *et. al.* with an additional assumption: the high-excited levels thermalize much faster with the cooling agent than the other levels. Evaporation of particles, known to be important in the initial stages of the cooling process, is explicitly included.

I. INTRODUCTION

The development of cooling techniques has opened up the possibility of studying ultracold gases. In particular, the quantum degeneracy of bosons and fermions has been investigated. Basically there are two different processes which can be used to cool an atomic or molecular ensemble, evaporative cooling and sympathetic cooling. However, there are gases in which the interaction is too weak for evaporative cooling to work. In addition, this method fails for identical fermions at low temperatures because of the exclusion principle [1]. In this case, sympathetic cooling can be used as an alternative to evaporative cooling: a cold gas is in thermal contact with another gas to be cooled. The first application of the sympathetic cooling method was in the context of the cooling of charged particles [2] and later it was extended to neutral atoms [3]. Only recently has it been applied in the context of ultracold atoms. In particular, quantum degeneracy of bosons and fermions has been achieved by thermalization of atoms of the same species in different internal states [4], between two isotopes of the same species [5], and finally between atoms of different species [6–9]. Moreover, in contrast to evaporative cooling, sympathetic cooling does not lead to a significant loss of either the cooled gas or the cooling agent.

From the theoretical point of view, classical and quantum models have been derived to describe the dynamics of thermalization of one-species and two-species systems. In the classical Boltzmann regime, an analytical formula for the evolution of the temperature in a mixture of nonequal mass atoms and equal mass atoms has been derived in Refs. [7,10]. A quantum mechanical description is given in terms of Boltzmann equations [11] and master equations [12–15]. In Ref. [11], sympathetic cooling of an atomic Fermi gas by a Bose gas is studied. In particular, the equilibrium temperature and the relaxation dynamics are obtained. In Ref. [12], for the first time, a master equation describing the sympathetic cooling of a system A is derived, treating the cooling agent B as a heat bath at a fixed temperature. More recently, by means of decoherence and ergodicity arguments a rate equation for sympathetic cooling of harmonically trapped bosons or fermions is obtained [13]. In Refs. [14,15] a detailed quantum kinetic master equation, which couples the dynamics of a trapped condensate to the vapor of noncondensate particles, is obtained along with information about the growth of a Bose-Einstein condensate. In Refs. [16–24], one can find more information about the growth of the condensate.

The purpose of this work is to describe the sympathetic cooling process of a total-number-conserving system in terms of particles in the ground state and elementary excitations. We use the particle-number-conserving Girardeau-Arnowitt formalism [25] and the sympathetic cooling model developed by M. Lewenstein et al. [12] with the additional assumption, that, the highly excited levels of the trap thermalize much faster with the cooling agent than the other levels (or in other words, the highly excited levels of the trap are in thermal equilibrium with the cooling agent). The advantage of the Girardeau-Arnowitt formalism is that it covers all cases ranging from a total absence of population in the ground state, $\mathsf{n}_0 = 0$, to a highly populated ground state $\mathsf{n}_0 \simeq N$. The assumption of the fast thermalization of the highly excited levels of the trap is related to the separation of the levels of the trap into two different bands, in the same way proposed by C.W. Gardiner et al. in Refs. [14,15], the *condensate* (B_C) band and the *noncondensate* band (B_{NC}), where the latter is in thermal equilibrium with the cooling agent. The condensate band includes all trap levels which are directly influenced by the presence of the condensate. The noncondensate band contains energy levels that are sufficiently high for the interaction with

---

*e-mail: andreans@mpipks-dresden.mpg.de

the condensate to be negligible. As we will show, the master equation for sympathetic cooling can be written in terms of two processes: (a) Creation and annihilation of particles in the ground state and (b) Creation and annihilation of quasiparticles. From this master equation we derive a rate equation for this number-conserving system under sympathetic cooling which includes the growth, scattering and loss (sympathetic evaporation) processes. This result is very important, specially now that sympathetic cooling is used as an alternative cooling method. In addition, this rate equation is more general than the one obtained by Gardiner *et. al.* in refs. [14,15], which is only valid in the regime of a highly populated condensate [26].

The paper is organized as follows: In section II the description of the sympathetic cooling model is given. In section III the master equation describing the dynamics of a number-conserving system in the condensate band is derived. In subsections IV A, IV B, and IV C a rate equation describing the population growth, the effects of the scattered particles as well as the effects of the sympathetic evaporation (trap losses) is derived. Section V contains the conclusion.

II. DESCRIPTION OF THE SYSTEM

For the sake of completeness and notation we briefly review the main results of Refs. [12,13]. However, we introduce some modifications to the model presented in Ref. [12]: the cooling agent is a gas trapped in a harmonic potential, instead of a free gas, and the gas to be cooled is confined in an open trap. This situation is closer to the actual experimental conditions.

A system A of N_A bosons is subject to sympathetic cooling due to its interaction with system B. The cooling agent of N_B atoms is in thermal equilibrium at temperature T_B. The single-particle states of system B are then described by harmonic oscillator wave functions with energy $\epsilon_{\vec{\ell}} = \hbar\omega_B(\ell_x + \ell_y + \ell_z)$, where ω_B is the trap frequency and $\vec{\ell} = (\ell_x, \ell_y, \ell_z)$ are the quantum numbers related to the bath. The corresponding Hamiltonian can be written

$$H_B = \sum_{\vec{\ell}} \hbar\omega_B(\ell_x + \ell_y + \ell_z) b^\dagger_{\vec{\ell}} b_{\vec{\ell}},$$

with the creation and annihilation operators for the bath $b^\dagger_{\vec{\ell}}$ and $b_{\vec{\ell}}$, respectively.

System A is assumed to be confined in an open trap. For energies ϵ much smaller than the trap depth ϵ_t, the trap can be approximated by a harmonic potential with trap frequency ω_A. The single-particle states in the harmonic oscillator potential have the quantum numbers $\vec{n} = (n_x, n_y, n_z)$. The single–particle eigenfunctions are labelled $\psi_{\vec{n}}(\vec{x})$, the eigenvalues $\epsilon_{\vec{n}} = \hbar\omega_A(n_x + n_y + n_z)$. The associated creation and annihilation operators are denoted by $a^\dagger_{\vec{n}}$ and $a_{\vec{n}}$, respectively. If, on the other hand, the energy is much larger than the trap depth, the system A can be approximately described by free particles . In this case the states are described by plane waves with energy $\epsilon(\vec{\kappa}) = \hbar^2\vec{\kappa}^2/(2m)$, with $\vec{\kappa}$ denoting the wavevector of the particle.

$$H_A = \sum_{\vec{n}} U_{\vec{n}} a^\dagger_{\vec{n}} a_{\vec{n}} \tag{1}$$

where

$$U_{\vec{n}} = \begin{cases} \epsilon_{\vec{n}} & \text{if } \epsilon \ll \epsilon_t \\ \epsilon(\vec{\kappa}) & \text{if } \epsilon \gg \epsilon_t \end{cases}$$

when $\epsilon \approx \epsilon_t$, $U_{\vec{n}}$ cannot be determined in a simple way.

The two-body interaction between the bath and the particles can be written in the form

$$V_{A-B} = \sum_{\vec{n},\vec{n'},\vec{\ell},\vec{\ell'}} \gamma_{\vec{n},\vec{n'},\vec{\ell},\vec{\ell'}} a^\dagger_{\vec{p}} a_{\vec{p'}} b^\dagger_{\vec{\ell}} b_{\vec{\ell'}}.$$

For simplicity, we assume a point-like interaction characterized by a scattering length a. We are interested in scattering processes between the bath and tightly bound particles, and processes in which low-energy particles of the system are scattered into the continuum through the interaction with the bath. In the first case, the matrix elements of the two-body interaction have the form

$$\gamma_{\vec{n},\vec{n'},\vec{\ell},\vec{\ell'}} = \frac{4\pi\hbar^2 a}{2\mu_{AB}} \int d\vec{x} \psi^*_{\vec{n}}(\vec{x}) \psi_{\vec{n'}}(\vec{x}) \psi^*_{\vec{\ell}}(\vec{x}) \psi_{\vec{\ell'}}(\vec{x}). \tag{2}$$

Here, $\mu_{AB} = m_A m_B/(m_A + m_B)$ denotes the reduced mass. For scattering processes into the continuum, the matrix element can be written as

$$\gamma_{\vec{n},\vec{\ell},\vec{\ell'}}(\vec{\kappa}) = \frac{4\pi\hbar^2 a}{2\mu_{AB}(2\pi)^{3/2}} \int d\vec{x} \quad e^{-i\vec{\kappa}\cdot\vec{x}} \psi_{\vec{n}}(\vec{x}) \psi^*_{\vec{\ell}}(\vec{x}) \psi_{\vec{\ell'}}(\vec{x}). \tag{3}$$

As derived in Ref. [12], the master equation for the time dependence of the reduced density matrix $\rho_A(t)$, obtained by tracing the total density matrix over the bath, has the form

$$\frac{\mathrm{d}\rho_{\mathrm{A}}(\mathrm{t})}{\mathrm{dt}} = -\frac{i}{\hbar}\left[H_A + V'_{A-A}, \rho_A(t)\right] + \mathcal{L}\rho_A(t), \tag{4}$$

where V'_{A-A} describes the mutual interaction of the atoms in system A. The interaction includes the two-body interaction V_{A-A} and, in addition, terms corresponding to shifts produced by the elimination of the bath in the master equation. It is assumed that the interaction V_{A-A} is weak compared to V_{A-B}, i.e. the thermalization of the system is solely determined by the interaction with the bath. The equation above is obtained under the the following assumptions: (i) Markov:

the correlation time for the interaction between systems A and B is much shorter than the cooling time, and (ii) rotating–wave approximation: terms rotating at multiples of the trap frequency are neglected.

It was shown in Ref. [13] that if the number N_B of particles in the bath is very large, decoherence acts very quickly compared to the equilibration time and reduces the density matrix to diagonal form. If only scattering processes involving tightly bound atoms of system A are involved, the Liouvillian in Eq. (4) becomes

$$\mathcal{L}\rho_{\mathcal{A}} = \sum_{\vec{m}\neq\vec{n}} \Gamma^{\vec{m},\vec{n}} \Big(2a^{\dagger}_{\vec{m}} a_{\vec{n}} \rho_A(t) a^{\dagger}_{\vec{n}} a_{\vec{m}} - a^{\dagger}_{\vec{n}} a_{\vec{m}} a^{\dagger}_{\vec{m}} a_{\vec{n}} \rho_A(t) - \rho_A(t) a^{\dagger}_{\vec{n}} a_{\vec{m}} a^{\dagger}_{\vec{m}} a_{\vec{n}} \Big), \quad (5)$$

with the rate coefficients

$$\Gamma^{\vec{m},\vec{n}} = \frac{1}{2\hbar^2} \int_{-\infty}^{\infty} \mathrm{d}\tau \sum_{\vec{\ell},\vec{\ell'}} \gamma_{\vec{n},\vec{m},\vec{\ell},\vec{\ell'}}\ \gamma_{\vec{m},\vec{n},\vec{\ell'},\vec{\ell}} \times \mathsf{n}_{\vec{\ell}} [\mathsf{n}_{\vec{\ell'}} + 1] \exp\left[i \left(\Delta\epsilon_A + \Delta\epsilon_B \right) \tau / \hbar \right] \quad (6)$$

where $\mathsf{n}_{\vec{\ell}}$ is the average occupation number of the heat bath oscillator. The energy difference for the system and the bath is given by $\Delta\epsilon_A = \epsilon_{\vec{n}} - \epsilon_{\vec{m}}$ and $\Delta\epsilon_B = \epsilon_{\vec{\ell}} - \epsilon_{\vec{\ell'}}$, respectively. If particles are scattered into the continuum, the rate coefficients have similar form but with the matrix element defined in Eq. (3). More details about the evaluation of the rate coefficients are given in Refs. [12,13].

III. MASTER EQUATION FOR THE CONDENSATE BAND

In this section, a master equation describing the dynamics of the condensate band is derived. Basically, we extend the formalism developed by Gardiner et al. [14] to the case of two distinct gases. The trap-levels of the system A are grouped in two bands: the *condensate* band (B_C) and the band of *noncondensed* particles (B_{NC}), where the latter is in thermal equilibrium with the cooling agent system B (see Figure 1).



FIG. 1. Distribution of the trap levels in the condensate B_C and noncondensate B_{NC} bands.

The condensate band B_C includes the ground state and all excited levels in the trap which are directly influenced by the presence of the condensate. The noncondensate band B_{NC} is composed by the highly excited levels, which include also the continuum states for $\epsilon > \epsilon_t$. The following assumptions are used in order to derive a master equation for the condensate band from Eq. (5):

(i) We neglect the correlations between the particles in the condensate band and the particles in the noncondensate band. Therefore, we assume that the complete density matrix of the system A can be written as a direct product between the density of the condensate band and the density of the noncondensate band, i.e., $\rho_A = \rho_C \otimes \rho_{NC}$. Since the main interest is in the dynamics of the condensate band, the noncondensate band is eliminated from the description by tracing out the noncondensate variable, i.e., $\rho_C = \mathrm{Tr}_{\mathrm{NC}}(\rho_{\mathrm{A}})$.

(ii) The density matrix of the noncondensate band ρ_{NC} is considered in thermal equilibrium with the cooling agent at temperature T_B. The levels of this band thermalize much faster with the cooling agent than the levels of the condensate band. Due to this assumption, we will completely neglect collisions between particles of the noncondensate band and particles of the cooling agent.

Applying these assumptions to Eq.(5), a master equation for the condensate band can be derived,

$$\frac{d\rho_C(t)}{dt} = \sum_{\substack{\vec{m}\in \mathrm{B_{NC}} \\ \vec{n}\in \mathrm{B_C}}} \Gamma^{\vec{n},\vec{m}}\ \langle N_{\vec{m}} \rangle_{\mathsf{n}_0} \Big(2a^{\dagger}_{\vec{n}} \rho_C(t) a_{\vec{n}} - a_n a^{\dagger}_n \rho_C(t) - \rho_C(t) a_{\vec{n}} a^{\dagger}_{\vec{n}} \Big) \quad (7)$$

$$+ \sum_{\substack{\vec{m}\in \mathrm{B_{NC}} \\ \vec{n}\in \mathrm{B_C}}} \Gamma^{\vec{m},\vec{n}}\ \langle N_{\vec{m}} + 1 \rangle_{\mathsf{n}_0} \Big(2a_{\vec{n}} \rho_C(t) a^{\dagger}_{\vec{n}} - a^{\dagger}_{\vec{n}} a_{\vec{n}} \rho_C(t) - \rho_C(t) a^{\dagger}_{\vec{n}} a_{\vec{n}} \Big) \quad (8)$$

$$+ \sum_{\substack{\vec{m}\neq\vec{n} \\ \vec{n},\vec{m}\in \mathrm{B_C}}} \Gamma^{\vec{m},\vec{n}} \Big(2a^{\dagger}_{\vec{m}} a_{\vec{n}} \rho_C(t) a^{\dagger}_{\vec{n}} a_{\vec{m}} - a^{\dagger}_{\vec{n}} a_{\vec{m}} a^{\dagger}_{\vec{m}} a_{\vec{n}} \rho_C(t) - \rho_C(t) a^{\dagger}_{\vec{n}} a_{\vec{m}} a^{\dagger}_{\vec{m}} a_{\vec{n}} \Big) \quad (9)$$

$$+ \sum_{\substack{\vec{m} \neq \vec{n} \\ \vec{n}, \vec{m} \in \mathrm{B_C}}} \Gamma^{\vec{n},\vec{m}} \Big(2a^{\dagger}_{\vec{n}} a_{\vec{m}} \rho_C(t) a^{\dagger}_{\vec{m}} a_{\vec{n}} - a^{\dagger}_{\vec{m}} a_{\vec{n}} a^{\dagger}_{\vec{n}} a_{\vec{m}} \rho_C(t) - \rho_C(t) a^{\dagger}_{\vec{m}} a_{\vec{n}} a^{\dagger}_{\vec{n}} a_{\vec{m}} \Big) \qquad (10)$$

where $\Gamma^{\vec{m},\vec{n}} = \Gamma^{\vec{n},\vec{m}} e^{-\beta(\epsilon_{\vec{m}} - \epsilon_{\vec{n}})}$ and $\langle N_{\vec{m}} \rangle_{\mathsf{n}_0} \equiv \mathrm{Tr}(\mathrm{a}^{\dagger}_{\vec{\mathrm{m}}} \mathrm{a}_{\vec{\mathrm{m}}} \rho_{\mathrm{NC}})_{\mathsf{n}_0}$ is the expectation value of the particle number operator for state $\vec{m}$, given that there are n_0 bosons in the ground state. Since the terms with $\vec{m} = \vec{n}$ cancel, this case is already excluded from the summation. The first two terms, Eqs.(7,8), account for collisions between particles in the condensate band and in the noncondensate band, and are illustrated in Fig.2. The loss of particles from the trap, due to their interaction with the cooling agent atoms are also included in Eq.(8). The existence of such a process is related to the fact that the trap is open, hence the atoms with higher energy than the trap energy can indeed escape. In the present model these particles are considered to escape to continuum states (non-bound states). In this case the single-particle states are described by plane waves. Evaporation is illustrated in Fig.3. The following two terms, Eqs.(9,10), describe collisions only between particles in the condensate band, which are shown in Fig.4.



FIG. 2. Collisions between particles in the condensate band B_C and in the noncondensate band B_{NC}



FIG. 3. Loss of particles from the trap, due to their interaction with the cooling agent atoms.



FIG. 4. Collisions between particles in the condensate band B_C.

Eqs. (7-10) give an accurate treatment of the internal dynamics of the condensate band, describing how the condensate forms and how the particles are scattered by collisions. Effects of elementary excitations of the condensate, representing the "thermal cloud", do not appear explicitly, since they are hidden in the single-particle operators $a_{\vec{n}}$ and $a^{\dagger}_{\vec{n}}$ for $\vec{n} \in \mathrm{B_C}$. In the limit of a highly condensed ground state, i.e. $\mathsf{n}_0 \simeq N$, the effects of the condensate excitations can be totally neglected [15]. During the process of condensate formation, however, their effects play a crucial role.

To analyse the role of elementary excitations, we resort to a formalism of Girardeau and Arnowitt [25], which conserves the number of particles. The method is based on the annihilation $\hat{\beta}_0$ and creation $\hat{\beta}^{\dagger}_0$ operators of one particle in the ground state, defined as $\hat{\beta}_0 = (\hat{N}_0 + 1)^{-1/2} \hat{a}_0$ and $\hat{\beta}^{\dagger}_0 = \hat{a}^{\dagger}_0 (\hat{N}_0 + 1)^{-1/2}$, and in the phonon operators $\hat{\beta}_{\vec{n}}$, $\hat{\beta}^{\dagger}_{\vec{n}}$ defined as $\hat{\beta}_{\vec{n}} = \hat{\beta}^{\dagger}_0 \hat{a}_{\vec{n}}$ and $\hat{\beta}^{\dagger}_{\vec{n}} = \hat{a}^{\dagger}_{\vec{n}} \hat{\beta}_0$, where $\hat{N}_0$ is the number operator of particles in the ground-state, $a_{\vec{n}}$ and $a^{\dagger}_{\vec{n}}$ are usual annihilation and creation operators for the trap level $\vec{n}$. These operators obey the Bose commutation relations, $[\hat{\beta}_{\vec{n}}, \hat{\beta}^{\dagger}_{\vec{n}'}] = \delta_{\vec{n},\vec{n}'}$. The phonons operators can be written in terms of the creation and annihilation quasiparticles operators: $\hat{b}_{\vec{n}}$ and $\hat{b}^{\dagger}_{\vec{n}}$, i.e., $\hat{\beta}_{\vec{n}} = u_{\vec{n}} \hat{b}_{\vec{n}} + v_{\vec{n}} \hat{b}^{\dagger}_{\vec{n}}$ and $\hat{\beta}^{\dagger}_{\vec{n}} = u^{*}_{\vec{n}} \hat{b}^{\dagger}_{\vec{n}} + v^{*}_{\vec{n}} \hat{b}_{\vec{n}}$, with $|u_{\vec{n}}|^2 - |v_{\vec{n}}|^2 = 1$. The formalism of Girardeau and Arnowitt is more general than the one developed by Gardiner [26], since it covers all cases ranging from a total absence of population in the ground state $\mathsf{n}_0 = 0$ to a highly populated ground state $\mathsf{n}_0 \simeq N$. A detailed comparison of both formalisms can be found in Ref. [27].

We replace the operators $\hat{a}_0, \hat{a}^{\dagger}_0, \hat{a}_{\vec{n}}, \hat{a}^{\dagger}_{\vec{n}}$ in Eqs. (7-10) by the particle-number-conserving Girardeau-Arnowitt operators $\hat{\beta}_0, \hat{\beta}^{\dagger}_0, \hat{\beta}_{\vec{n}}, \hat{\beta}^{\dagger}_{\vec{n}}$. Then, the action of the opertators on the density matrix $\rho_C(t)$ is computed. Since the decoherence time is very fast (see discussion in Ref. [13]), only the diagonal terms of the reduced density matrix ρ_C contribute to the dynamics. Hence, only diagonal terms will be considered. In this way, we obtain a master equation for the diagonal elements $P(\mathsf{n}_0, \vec{n}) = \langle \mathsf{n}_0, \vec{n} | \rho_C(t) | \mathsf{n}_0, \vec{n} \rangle$, with $\vec{n} \equiv (\mathsf{n}_1, ..., \mathsf{n}_m, \mathsf{n}_n, ...)$ denoting the set of occupation number for all quasiparticles:

$$\frac{d}{dt}P(\mathsf{n}_0,\vec{n},t) = \frac{d}{dt}P_{\text{growth}}(\mathsf{n}_0,\vec{n},t) + \frac{d}{dt}P_{\text{scatt}}(\mathsf{n}_0,\vec{n},t) + \frac{d}{dt}P_{\text{evap}}(\mathsf{n}_0,\vec{n},t). \tag{11}$$

The rate $dP_{\text{growth}}(\mathsf{n}_0,\vec{n},t)/dt$ represents the contribution of the condensate itself and the quasiparticles for the growth of the condensate. The number of particles in the condensate band changes. $dP_{\text{scatt}}(\mathsf{n}_0,\vec{n},t)/dt$ describes scattered particles in the condensate band. During the scattering process the number of particles in the condensate band does not change. Finally, $dP_{\text{evap}}((\mathsf{n}_0,\vec{n},t)/dt$ includes the evaporation of particles with energies larger than the trap depth.

IV. RATE EQUATIONS FOR THE CONDENSATE BAND

A. Growth

Consider the diagonal elements of the Eqs. (7–8) for $\vec{n}=0$ and $\vec{n}\neq 0$ with the replacements described in the previous section, we arrive at a master equation for the occupation probability of the condensate band

$$\frac{d}{dt}P_{\text{growth}}(\mathsf{n}_0,\vec{n},t) = R_C + R_Q, \tag{12}$$

where R_C represents the growth rate of the condensate itself

$$\begin{aligned} R_C = {} & 2\mathsf{n}_0\Gamma^+(\mathsf{n}_0-1)P(\mathsf{n}_0-1,\vec{n}) - 2(\mathsf{n}_0+1)\Gamma^+(\mathsf{n}_0)P(\mathsf{n}_0,\vec{n}) \\ & +2(\mathsf{n}_0+1)\Gamma^-(\mathsf{n}_0+1)P(\mathsf{n}_0+1,\vec{n}) - 2\mathsf{n}_0\Gamma^-(\mathsf{n}_0)P(\mathsf{n}_0,\vec{n}). \end{aligned} \tag{13}$$

The coefficients $\Gamma^+(\mathsf{n}_0)$ and $\Gamma^-(\mathsf{n}_0)$ carry information about particles entering and leaving the condensate

$$\begin{aligned} \Gamma^+(\mathsf{n}_0) &\equiv \sum_{\vec{m}\neq 0} \Gamma^{\vec{0},\vec{m}} \langle N_{\vec{m}}\rangle_{\mathsf{n}_0}, \\ \Gamma^-(\mathsf{n}_0) &\equiv \sum_{\vec{m}\neq 0} \Gamma^{\vec{m},\vec{0}} \langle N_{\vec{m}}+1\rangle_{\mathsf{n}_0}. \end{aligned}$$

The rate R_Q takes into account the contribution of the elementary excitations

$$\begin{aligned} R_Q = {} & 2\sum_{\vec{n}} \mathsf{n}_{\vec{n}}\Gamma^{++}_{\vec{n}}(\mathsf{n}_0-1,\vec{n}-\vec{e}_{\vec{n}})P(\mathsf{n}_0-1,\vec{n}-\vec{e}_{\vec{n}}) \\ & -2\sum_{\vec{n}}(\mathsf{n}_{\vec{n}}+1)\Gamma^{++}_{\vec{n}}(\mathsf{n}_0,\vec{n})P(\mathsf{n}_0,\vec{n}) \\ & +2\sum_{\vec{n}}(\mathsf{n}_{\vec{n}}+1)\Gamma^{--}_{\vec{n}}(\mathsf{n}_0+1,\vec{n}+\vec{e_{\vec{n}}})P(\mathsf{n}_0+1,\vec{n}+\vec{e_{\vec{n}}}) \\ & -2\sum_{\vec{n}} \mathsf{n}_{\vec{n}}\Gamma^{--}_{\vec{n}}P(\mathsf{n}_0,\vec{n}) \\ & +2\sum_{\vec{n}}(\mathsf{n}_{\vec{n}}+1)\Gamma^{+-}_{\vec{n}}(\mathsf{n}_0-1,\vec{n}+\vec{e_{\vec{n}}})P(\mathsf{n}_0-1,\vec{n}+\vec{e_{\vec{n}}}) \\ & -2\sum_{\vec{n}} \mathsf{n}_{\vec{n}}\Gamma^{+-}_{\vec{n}}(\mathsf{n}_0)P(\mathsf{n}_0,\vec{n}) \\ & +2\sum_{\vec{n}} \mathsf{n}_{\vec{n}}\Gamma^{-+}_{n}(\mathsf{n}_0+1,\vec{n}-\vec{e_{\vec{n}}})P(\mathsf{n}_0+1,\vec{n}-\vec{e_{\vec{n}}}) \\ & -2\sum_{\vec{n}}(\mathsf{n}_{\vec{n}}+1)\Gamma^{-+}_{\vec{n}}(\mathsf{n}_0,\vec{n})P(\mathsf{n}_0,\vec{n}), \end{aligned} \tag{14}$$

with $\vec{e_n} \equiv (...,0,0,1,0,0,...)$. The coefficients $\Gamma^{++}_{\vec{n}}(\mathsf{n}_0,\vec{n})$, $\Gamma^{--}_{\vec{n}}(\mathsf{n}_0,\vec{n})$, $\Gamma^{+-}_{\vec{n}}(\mathsf{n}_0,\vec{n})$ and $\Gamma^{-+}_{\vec{n}}(\mathsf{n}_0,\vec{n})$ are defined as

$$\begin{aligned} \Gamma^{++}_{\vec{n}}(\mathsf{n}_0,\vec{n}) &\equiv \sum_{\vec{m}\neq\vec{n}} \Gamma^{\vec{n},\vec{m}}\langle N_{\vec{m}}\rangle_{\mathsf{n}_0}|u_{\vec{n}}|^2, \\ \Gamma^{--}_{\vec{n}}(\mathsf{n}_0,\vec{n}) &\equiv \sum_{\vec{m}\neq\vec{n}} \Gamma^{\vec{m},\vec{n}}\langle N_{\vec{m}}+1\rangle_{\mathsf{n}_0}|u_{\vec{n}}|^2, \\ \Gamma^{+-}_{\vec{n}}(\mathsf{n}_0,\vec{n}) &\equiv \sum_{\vec{m}\neq\vec{n}} \Gamma^{\vec{n},\vec{m}}\langle N_{\vec{m}}\rangle_{\mathsf{n}_0}|v_{\vec{n}}|^2, \\ \Gamma^{-+}_{\vec{n}}(\mathsf{n}_0,\vec{n}) &\equiv \sum_{\vec{m}\neq\vec{n}} \Gamma^{\vec{m},\vec{n}}\langle N_{\vec{m}}+1\rangle_{\mathsf{n}_0}|v_{\vec{n}}|^2, \end{aligned}$$

where $\vec{m}\in B_{NC}$ and $\vec{n}\in B_C$. These coefficients describe processes of creation and annihilation of phonons inside the condensate.

In order to obtain the dynamics of the population in the condensate band, the mean number of particles is evaluated. We will start with the contribution of the condensate mode itself. In Eq.(13) we use the thermodynamic relation between the rate coefficients

$$\Gamma^+(\mathsf{n}_0) = \Gamma^-(\mathsf{n}_0)e^{\beta(\epsilon_0-\mu_0)}$$

in combination with the factorization assumption $P(\mathsf{n}_0,\vec{n}) = P(\mathsf{n}_0)\otimes P(\vec{n}) = P(\mathsf{n}_0)\ldots\otimes P(\mathsf{n}_n)\otimes\ldots$. Here $\beta = 1/k_BT_B$, ϵ_0 is the energy of the ground state and $\mu_0 \equiv \mu(\mathsf{n}_0)$ the chemical potential for n_0 particles in the ground state. For the rate of change of the condensate number one then finds

$$\begin{aligned} \frac{d\mathsf{n}_0}{dt}|_{\text{growth}} &= 2\Big((\mathsf{n}_0+1)\Gamma^+(\mathsf{n}_0) - \mathsf{n}_0\Gamma^-(\mathsf{n}_0)\Big) \\ &= 2\Gamma^+(\mathsf{n}_0)\Big(\mathsf{n}_0(1-e^{-\beta(\mu_0-\epsilon_0)})+1\Big). \end{aligned} \tag{15}$$

In the limit of $\mathsf{n}_0 \simeq N$, with $N \gg 1$, the equation above can be approximated by

$$\frac{dN}{dt} \approx 2\Gamma^+(N)N\Big((1-e^{\beta(\mu_0-\epsilon_0)})\Big), \tag{16}$$

where the coefficient $\Gamma^+(N)$ contains information about the sympathetic cooling process between system A and cooling agent B.

If we apply the formalism described here to a gas of only one species, we find a rate equation for the growth of the condensate which has a form similar to the one obtained by Gardiner and coworkers based on [26] (see Eq. (23) in Ref. [15] and Eq.(168) in Ref. [14]). However, due to the properties of the Giradeau-Arnowitt formalism, the rate equation in our case is not limited only to highly populated condensates but also includes condensates of arbitrary particle number.

We now explore the contribution of the elementary excitations for the growth process. We use the thermodynamic relation between the coefficients

$$\Gamma^{++}_{\vec n}(\mathsf{n}_0,\vec n) = \Gamma^{--}_{\vec n}(\mathsf{n}_0,\vec n) e^{\beta(\epsilon_{\vec n}-\mu_0)},$$

$$\Gamma^{+-}_{\vec n}(\mathsf{n}_0,\vec n) = \Gamma^{-+}_{\vec n}(\mathsf{n}_0,\vec n) e^{\beta(\epsilon_{\vec n}-\mu_0)},$$

and factorize all correlations i.e., $P(\mathsf{n}_0,\vec n) = P(\mathsf{n}_0)\otimes ...P(\mathsf{n}_n)...$ in Eq.(14). Thus, the rate equation for the elementary excitations is given by

$$\frac{d\mathsf{n}_{\vec n}}{dt}|_{\text{growth}} = 2\sum_{\vec n}\Gamma^{++}_{\vec n}(\mathsf{n}_0,\vec n)\Bigg((1-e^{\beta(\mu_0-\epsilon_{\vec n})})\mathsf{n}_{\vec n}+1\Bigg)$$
$$-2\sum_{\vec n}\Gamma^{+-}_{\vec n}(\mathsf{n}_0,\vec n)\Bigg((1-e^{\beta(\mu_0-\epsilon_{\vec n})})\mathsf{n}_{\vec n}+1\Bigg). \quad (17)$$

A simplified form of Eq.(17) can be derived by using the relation $\Gamma^{+-}_{\vec n}(\mathsf{n}_0,\vec n) = \Gamma^{++}_{\vec n}(\mathsf{n}_0,\vec n) - \Gamma^{+}_{\vec n}(\mathsf{n}_0,\vec n)$,

$$\frac{d\mathsf{n}_{\vec n}}{dt}|_{\text{growth}} = 2\sum_{\vec n}\Gamma^{+}_{\vec n}(\mathsf{n}_0,\vec n)\Bigg((1-e^{\beta(\mu_0-\epsilon_{\vec n})})\mathsf{n}_{\vec n}+1\Bigg), \quad (18)$$

with $\Gamma^{+}_{\vec n}(\mathsf{n}_0,\vec n) = \sum_{\vec m}\Gamma^{\vec n,\vec m}\langle N_{\vec m}\rangle_{\mathsf{n}_0}$.

The total growth equation can be obtained from the contribution of the condensate mode itself Eq.(15) and the elementary excitations Eq.(18).

B. Scattering

In this subsection the effects of the scattering processes of atoms inside the condensate band are considered, as described by Eqs. (9−10). During the scattering process the number of particles in the condensate band does not change.

We use the same analytical procedure described in the previous section to derive a rate equation related to the scattering process. However, we will omit details of this calculation, since the expressions can be rather large. Thus, the corresponding scattering rate equation for $\mathsf{n}_{\vec n} = \langle a^\dagger_{\vec n} a_{\vec n}\rangle$ has the form

$$\frac{d\mathsf{n}_{\vec n}}{dt}|_{\text{scatt}} = 2\sum_{\vec n\neq 0}\Gamma^{u+}_{\vec n}\mathsf{n}_0[e^{-\beta(\epsilon_{\vec n}-\epsilon_0)} - \mathsf{n}_{\vec n}(1-e^{-\beta(\epsilon_{\vec n}-\epsilon_0)})]$$
$$+2\sum_{\vec n\neq 0}\Gamma^{v+}_{\vec n}\mathsf{n}_0[\mathsf{n}_{\vec n}(1-e^{-\beta(\epsilon_{\vec n}-\epsilon_0)})+1]$$
$$+2\sum_{\vec m\neq\vec n\neq 0}\Gamma^{uu+-}_{\vec m,\vec n}[\mathsf{n}_{\vec m}(\mathsf{n}_{\vec n}+1)$$
$$-\mathsf{n}_{\vec n}(\mathsf{n}_{\vec m}+1)e^{-\beta(\epsilon_{\vec m}-\epsilon_{\vec n})}]$$
$$+2\sum_{\vec m\neq\vec n\neq 0}\Gamma^{vv+-}_{\vec m,\vec n}[-\mathsf{n}_{\vec n}(\mathsf{n}_{\vec m}+1)$$
$$+\mathsf{n}_{\vec m}(\mathsf{n}_{\vec n}+1)e^{-\beta(\epsilon_{\vec m}-\epsilon_{\vec n})}]$$
$$+2\sum_{\vec m\neq\vec n\neq 0}\Gamma^{uv++}_{\vec m,\vec n}[(\mathsf{n}_{\vec m}+1)(\mathsf{n}_{\vec n}+1)$$
$$-\mathsf{n}_{\vec n}\mathsf{n}_{\vec m}e^{-\beta(\epsilon_{\vec m}-\epsilon_{\vec n})}]$$
$$+2\sum_{\vec m\neq\vec n\neq 0}\Gamma^{vu++}_{\vec m,\vec n}[-\mathsf{n}_{\vec n}\mathsf{n}_{\vec m}$$
$$+(\mathsf{n}_{\vec m}+1)(\mathsf{n}_{\vec n}+1)e^{-\beta(\epsilon_{\vec m}-\epsilon_{\vec n})}], \quad (19)$$

with

$$\Gamma^{u+}_{\vec n}(\mathsf{n}_0,\vec n) \equiv \Gamma^{0,\vec n}_{\vec n,0}|u_{\vec n}|^2,$$
$$\Gamma^{v+}_{\vec n}(\mathsf{n}_0,\vec n) \equiv \Gamma^{0,\vec n}_{\vec n,0}|v_{\vec n}|^2,$$
$$\Gamma^{uu+-}_{\vec n,\vec m}(\mathsf{n}_0,\vec n) \equiv \Gamma^{\vec n,\vec m}|u_{\vec n}|^2|u_{\vec m}|^2,$$
$$\Gamma^{vv+-}_{\vec n,\vec m}(\mathsf{n}_0,\vec n) \equiv \Gamma^{\vec n,\vec m}|v_{\vec n}|^2|v_{\vec m}|^2,$$
$$\Gamma^{uv++}_{\vec n,\vec m}(\mathsf{n}_0,\vec n) \equiv \Gamma^{\vec n,\vec m}|u_{\vec n}|^2|v_{\vec m}|^2,$$
$$\Gamma^{vu++}_{\vec n,\vec m}(\mathsf{n}_0,\vec n) \equiv \Gamma^{\vec n,\vec m}|v_{\vec n}|^2|u_{\vec m}|^2. \quad (20)$$

The first two terms of Eq.(19) account for the scattering between the particles in the ground state and the quasiparticles of the $\vec n$-level. The other terms describe the scattering between quasiparticles of different levels.

C. Evaporation

We now discuss the evaporation of particles from the trap. Atoms can escape from the condensate band of the trap to unbound states. This loss of particles from the trap is induced by their interaction with the cooling agent. This process is particularly important in the initial stage of the cooling process [7,9]. To implement evaporation in the present model, particles are assumed to escape to continuum states (unbound states) inside the noncondensate band. Only Eq. (8) contributes to this process, since the rate coefficients of Eq. (7), which includes the overlap of the wave functions involved in the process, is completely negligible. Taking the diagonal elements of Eq. (8), P_{evap} is obtained

$$\frac{d}{dt}P_{\text{evap}}(\mathsf{n}_0,\vec n,t) = -2\mathsf{n}_0\gamma^-(\mathsf{n}_0)P(\mathsf{n}_0,\vec n)$$
$$+2(\mathsf{n}_0+1)\gamma^-(\mathsf{n}_0+1)P(\mathsf{n}_0+1,\vec n)$$
$$+2\sum_{\vec n\,\in\,\mathrm{B_C}}(\mathsf{n}_{\vec n}+1)\gamma^{--}_{\vec n}(\mathsf{n}_0+1,\vec n+\vec e_n)P(\mathsf{n}_0+1,\vec n+\vec e_n)$$
$$-2\sum_{\vec n\,\in\,\mathrm{B_C}}\mathsf{n}_{\vec n}\gamma^{--}(\mathsf{n}_0)P(\mathsf{n}_0,\vec n)$$

$$+2 \sum_{\vec{n} \,\in\, \mathrm{B_C}} \mathsf{n}_{\vec{n}} \gamma_{\vec{n}}^{-+}(\mathsf{n}_0+1, \vec{n}-\vec{e}_n) P(\mathsf{n}_0+1, \vec{n}-\vec{e}_n)$$
$$-2 \sum_{\vec{n} \,\in\, \mathrm{B_C}} (\mathsf{n}_{\vec{n}}+1)\gamma^{-+}(\mathsf{n}_0, \vec{n}) P(\mathsf{n}_0, \vec{n}), \tag{21}$$

where

$$\begin{aligned}
\gamma^{-}(\mathsf{n}_0, \vec{n}) &\equiv \sum_{\vec{m} \in \mathrm{B_{NC}^{(t)}}} \Gamma^{\vec{m},0}, \\
\gamma_{\vec{n}}^{--}(\mathsf{n}_0, \vec{n}) &\equiv \sum_{\vec{m} \in \mathrm{B_{NC}^{(t)}}} \Gamma^{\vec{m},\vec{n}} |u_{\vec{n}}|^2, \\
\gamma_{\vec{n}}^{-+}(\mathsf{n}_0, \vec{n}) &\equiv \sum_{\vec{m} \in \mathrm{B_{NC}^{(t)}}} \Gamma^{\vec{m},\vec{n}} |v_{\vec{n}}|^2 \\
&= \gamma_{\vec{n}}^{-+}(\mathsf{n}_0, \vec{n}) - \gamma_{\vec{n}}^{-},
\end{aligned} \tag{22}$$

with $\mathrm{B_{NC}^{(t)}}$ denoting the noncondensate band with energy $\epsilon_{\vec{m}} > \epsilon_t$. As mentioned at the end of section II, these rate coefficients $\Gamma^{\vec{m},0}$ and $\Gamma^{\vec{m},\vec{n}}$ have a different form from the ones which appear in the growth and scattering processes, since now $\vec{m}$ describes unbound states. The first line of Eq.(21) describes the evaporation of an atom out of the ground state while the others describe the evaporation of an atom out of the n-excited level of the condensate band.

Following the same procedure described in the previous subsections, a rate equation for the evaporation of particles from the ground state and the evaporation of quasiparticles in the condensate band is obtained:

$$\begin{aligned}
\frac{d\mathsf{n}_0}{dt}\Big|_{\mathrm{evap}} &= -2 \quad \gamma^{-}(\mathsf{n}_0, \vec{n}) \quad \mathsf{n}_0|_{\mathrm{evap}}, \\
\frac{d\mathsf{n}_{\vec{n}}}{dt}\Big|_{\mathrm{evap}} &= -2 \sum_{\vec{n}} \gamma_{\vec{n}}^{-}(\mathsf{n}_0, \vec{n}) \quad \mathsf{n}_{\vec{n}}|_{\mathrm{evap}},
\end{aligned} \tag{23}$$

with

$$\gamma_{\vec{n}}^{-} = \sum_{\vec{m} \in \mathrm{B_{NC}^{(t)}}} \Gamma_{\vec{n},\vec{m}}^{\vec{m},\vec{n}}. \tag{24}$$

V. CONCLUSION

The total rate equation describing a number-conserving population of the condensate band can be obtained from Eqs.(15,17,19,23)

$$\frac{d\mathsf{n}_{\vec{n}}}{dt} = \frac{d\mathsf{n}_{\vec{n}}}{dt}\Big|_{\mathrm{growth}} + \frac{d\mathsf{n}_{\vec{n}}}{dt}\Big|_{\mathrm{scatt}} + \frac{d\mathsf{n}_{\vec{n}}}{dt}\Big|_{\mathrm{evap}}, \tag{25}$$

where n_0 and $\mathsf{n}_{\vec{n}}$ denote, respectively, the population in the condensate and the population of the elementary excitations, which are commonly referred to as "thermal cloud". The equation above gives a complete description for the thermalization of a system A which is in thermal contact with a bath B. All the information about the dynamics of the thermalization process is contained in the coefficients Γ. Analytical and numerical evaluation of the coefficients for a specific system can be carried out along the lines described in Refs. [12,13].

The result obtained here can be applied to describe the dynamics of sympathetic cooling of a gas in thermal contact with a cooling agent, in terms of the population in the ground-state and elementary excitations. In particular, it can be aplied to the case where the cooling agent thermalize much faster compared to the thermalization of the system. The description remains valid for the quantum degenerate regime $T \ll T_c$, where T_c is the critical temperature of the gas.

The use of the Giradeau-Arnowitt method has opened up the possibility to describe sympathetic cooling in terms of particles in the ground state and elementary excitation during the whole cooling process, i.e., from the case of total absence of particles in the condensate ($\mathsf{n}_0 = 0$) to the case of a highly populated ground state ($\mathsf{n}_0 = N$). In addition, we include the effects of evaporation which are especially important during the initial stages of the cooling process. To our knowledge this is the first time that a formal complete number-conserving description of sympathetic cooling in terms of particles in the ground state and elementary excitations is given.

ACKNOWLEDGMENTS

ANS thanks fruitful discussions with M. da Mata, A.F.R. de Toledo Piza, M.O. da C. Pires, A. Mosk, M. Ameduri, H. Schomerus, A. Ozorio de Almeida and H.A. Weidenmüller. ANS gratefully acknowledges M. Weidemüller and M.C. Nemes for many discussions and help in preparing the manuscript.

---

[1] B. DeMarco *et al.*, Phys. Rev. Lett. **82**, 4208 (1999).
[2] D. J. Larson *et al.*, Phys. Rev. Lett. **57**, 70 (1986), and references therein.
[3] H. F. Hess *et al.*, Phys. Rev. Lett. **59**, 672 (1988); R. van Roijen *et al.*, Phys. Rev. Lett. **61**, 931 (1988);
[4] C. Myatt *et al.*, Phys. Rev. Lett. **78**, 586 (1997).
[5] A. G. Truscott *et al.*, Science **291**, 2570 (2001).
[6] F. Schreck *et al.*, Phys. Rev. Lett. **87**, 080403 (2001).
[7] A. Mosk *et al.*, Appl. Phys. **B 73**, 791 (2001).
[8] Z. Hadzibacbic *et al.*, Phys. Rev. Lett. **88**, 160401 (2002).
[9] M. Mudrich *et al.*, Phys. Rev. Lett. **88**, 253001 (2002).
[10] G. Delannoy *et al.*, Phys. Rev. **A 63**, 011402R (2001).
[11] W. Geist *et al.*, Phys. Rev. **A 59**, 1500 (1999).

[12] M. Lewenstein, J. I. Cirac, and P. Zoller, Phys. Rev. **A 51**, 4617 (1995).

[13] T. Papenbrock, A. N. Salgueiro, and H. A. Weidenmüller, Phys. Rev. **A 65**, 043601 (2002).

[14] C. W. Gardiner *et al.*, Phys. Rev. **A 58**, 536 (1998).

[15] C. W. Gardiner *et al.*, Phys. Rev. Lett. **79**, 1793 (1997).

[16] C. W. Gardiner *et al.*, Phys. Rev. Lett. **81**, 5266 (1998).

[17] C. W. Gardiner *et al.*, Phys. Rev. **A 55**, 2902 (1997).

[18] D. Jacksch *et al.*, Phys. Rev. **A 56**, 575 (1997).

[19] D. Jacksch *et al.*, Phys. Rev. **A 58**, 1450 (1998).

[20] C. W. Gardiner *et al.*, Phys. Rev. **A 61**, 033601 (2000).

[21] M. D. Lee *et al.*, Phys. Rev. **A 61**, 033601 (2000).

[22] M. J. Davis *et al.*, Phys. Rev. **A 62**, 063608 (2000).

[23] M. Köhl *et al.*, Phys. Rev. Lett. **88**, 080402 (2002).

[24] C. W. Gardiner *et al.*, J. Phys. **B 35**, 733 (2002).

[25] M. Girardeau and R. Arnowitt, Phys. Rev. **113**, 755 (1959).

[26] C. W. Gardiner, Phys. Rev. **A 56**, 1414 (1997).

[27] M. Girardeau, Phys. Rev. **A 58**, 775 (1998).